

STATS 217: Introduction to Stochastic Processes I

Lecture 13

Evolution of distributions

- Consider a DTMC $(X_n)_{n \geq 0}$ on S with transition matrix P .
- Suppose we start the chain from a random initial state distributed according to λ . We will use the notation $X_0 \sim \lambda$. This just means that

$$\mathbb{P}[X_0 = i] = \lambda_i \quad \forall i \in S.$$

- What is the distribution of X_1 ?
- More generally, what is the distribution of X_n ?

Evolution of distributions

For any $j \in S$, we have

$$\begin{aligned}\mathbb{P}[X_n = j] &= \sum_{i \in S} \mathbb{P}[X_0 = i \wedge X_n = j] \\ &= \sum_{i \in S} \mathbb{P}[X_n = j \mid X_0 = i] \mathbb{P}[X_0 = i] \\ &= \sum_{i \in S} \lambda_i \cdot p_{ij}^n \\ &= \sum_{i \in S} \lambda_i \cdot (P^n)_{ij} \\ &= (\lambda P^n)_j.\end{aligned}$$

Stationary distributions

- So, if $X_0 \sim \lambda$, then $X_n \sim \lambda P^n$.
- **A stationary distribution** for P is a probability distribution π on S satisfying

$$\pi P = \pi.$$

- Therefore, if π is a stationary distribution for P , then

$$X_0 \sim \pi \implies X_n \sim \pi \quad \forall n \geq 1.$$

Existence and uniqueness

- A Markov chain $(X_n)_{n \geq 0}$ on S with transition matrix P is said to be **irreducible** if all the states form a single communicating class.
- Recall that this means that for all $i, j \in S$, there exists some t (possibly depending on i and j) such that $(P^t)_{i,j} > 0$.
- Recall also that since S is finite, this means that all states in S are recurrent.
- Next time: Let P be the transition matrix of an irreducible Markov chain. Then, there exists a unique probability distribution π satisfying $\pi P = \pi$.

Example

Two state chain: $S = \{0, 1\}$ and for $p, q \in (0, 1]$,

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}$$

- Since $p, q > 0$, the chain is irreducible.
- By the theorem, there is a unique stationary distribution.
- By solving $\pi P = \pi$ and using that π is a probability distribution, we get (check!) the solution

$$\pi = \left(\frac{q}{p+q}, \frac{p}{p+q} \right).$$

Doubly-stochastic Markov chains

- Consider a DTMC on S with transition matrix P .
- We know that entries of each row of the transition matrix P sum to 1.
- Suppose also that the columns of P sum to 1. Then, P is said to be a **doubly-stochastic transition matrix**.
- For instance, last time, in our study of waiting times for patterns in coin tossing, we encountered the doubly-stochastic transition matrix

$$P = \begin{bmatrix} & HH & HT & TH & TT \\ HH & 1/2 & 1/2 & 0 & 0 \\ HT & 0 & 0 & 1/2 & 1/2 \\ TH & 1/2 & 1/2 & 0 & 0 \\ TT & 0 & 0 & 1/2 & 1/2 \end{bmatrix}.$$

- Problem 1, Homework 4: Let P be a doubly-stochastic transition matrix on the state space S . Then, the uniform distribution on S is a stationary distribution.

Detailed balance conditions

- Consider a DTMC on S with transition matrix P . Let μ be a probability distribution on S .
- We say that μ satisfies the **detailed balance conditions** with respect to P if

$$\mu_i P_{ij} = \mu_j P_{ji} \quad \forall i, j \in S.$$

- If μ satisfies the detailed balance conditions with respect to P , then μ is a stationary distribution for P .
- Indeed, for all $i \in S$,

$$(\mu P)_i = \sum_{j \in S} \mu_j P_{ji} = \sum_{j \in S} \mu_i P_{ij} = \mu_i \sum_{j \in S} P_{ij} = \mu_i.$$

Detailed balance conditions

- If π satisfies the detailed balance conditions with respect to P and $X_0 \sim \pi$, then

$$\begin{aligned}\mathbb{P}[X_0 = x_0, \dots, X_n = x_n] &= \pi_{x_0} P_{x_0, x_1} \cdots P_{x_{n-1}, x_n} \\ &= P_{x_1, x_0} \cdot \pi_{x_1} P_{x_1, x_2} \cdots P_{x_{n-1}, x_n} \\ &= P_{x_1, x_0} P_{x_2, x_1} \cdot \pi_{x_2} P_{x_2, x_3} \cdots P_{x_{n-1}, x_n} \\ &= \dots \\ &= \pi_{x_n} P_{x_n, x_{n-1}} \cdots P_{x_1, x_0} \\ &= \mathbb{P}[X_0 = x_n, \dots, X_n = x_0].\end{aligned}$$

- For this reason, such chains are also called **reversible**.
- In many interesting examples, the detailed balance conditions provide an efficient way of finding the stationary distribution.

Example: Random walk on a graph

- $G = (V, E)$ is a graph, where V is the set of vertices and E is the set of edges.
- For vertices $u \neq v \in V$, we say that $u \sim v$ if and only if there is an edge between u and v .
- For a vertex $u \in V$, $\deg(u)$ denotes the degree of u i.e. the number of vertices it is connected to.
- Note that $\sum_{u \in V} \deg(u) = 2|E|$.
- Recall that the transition matrix of the random walk is given by

$$P_{u,v} = \begin{cases} \frac{1}{\deg(u)} & \text{if } v \sim u \\ 0 & \text{otherwise.} \end{cases}$$

Example: Random walk on a graph

- Consider the distribution π where $\pi_u = \deg(u)/2|E|$.
- Then, π is a probability distribution.
- We claim that π satisfies the detailed balance conditions with respect to P .
- There are two cases. If $u \not\sim v$, then the condition is clearly satisfied since $P_{uv} = P_{vu} = 0$.
- If $u \sim v$, then

$$\pi_u P_{uv} = \frac{\deg(u)}{2|E|} \cdot \frac{1}{\deg(u)} = \frac{1}{2|E|} = \pi_v P_{vu}.$$

- Therefore, π is a stationary distribution for P . Note that P is irreducible if and only if the graph G is connected i.e., there is a path from any vertex to any other vertex, in which case, π is the unique stationary distribution.

Example: The Ehrenfest urn

The Ehrenfest urn. n balls are distributed among two urns, urn A and urn B . At each time, we select a ball uniformly at random and move it from its current urn to the other urn.

- Let X_t denote the number of balls in urn A at time t . Then, $(X_t)_{t \geq 0}$ is a DTMC on $\{0, \dots, n\}$ with transition matrix P given by

$$P_{jk} = \begin{cases} j/n & \text{if } k = j - 1 \\ (n - j)/n & \text{if } k = j + 1 \\ 0 & \text{otherwise.} \end{cases}$$

- Note that P is clearly irreducible.

Example: The Ehrenfest urn

- Let π be the distribution on $\{0, \dots, n\}$ given by

$$\pi_x = 2^{-n} \cdot \binom{n}{x}.$$

- By the binomial theorem, π is a probability distribution.
- **Exercise:** check that π satisfies the detailed balance condition with respect to P .
- Hence, π is the unique stationary distribution for P .